

Client
Gurugram
Pathkind Diagnostics Pvt. Ltd.
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

Processed By
Pathkind Diagnostics Pvt. Ltd.
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

| | |
|-----------------------------------|--|
| Name : Mr. CL103 | Billing Date : 07/07/2023 12:17:23 |
| Age : 35 Yrs | Sample Collected on : 10/07/2023 10:01:31 |
| Sex : Male | Sample Received on : 10/07/2023 11:02:13 |
| P. ID No. : P1000100012585 | Report Released on : 14/07/2023 18:20:27 |
| Accession No : 10002304641 | Barcode No. : 10002304641-01 |
| Referring Doctor : Self | Ref no. : |
| Referred By : | |

Report Status - Final

| Test Name | Result | Biological Ref. Interval | Unit |
|-----------|--------|--------------------------|------|
|-----------|--------|--------------------------|------|

BIOCHEMISTRY

| | | | |
|--|----------------|-------------|-------|
| Prostate Specific Antigen (PSA) Total | 25.00 H | 0.00 - 1.40 | ng/mL |
| <i>Sample: Serum</i> | | | |
| <i>Method: ECLIA</i> | | | |

Prostate Specific Antigen (PSA) Total

Prostate specific antigen (PSA Total) is a blood test that helps in the screening of prostate cancer. PSA is a protein produced by cancerous and noncancerous tissue in the prostate. This test is also used to monitor recurrence & response to treatment in known cases of prostate cancer. Many other conditions, such as an enlarged or inflamed prostate can also increase PSA levels. recommended test for detection of prostate cancer along with Digital Rectal Examination (DRE) in males above 50 years of age. Fa negative / positive results are observed in patients receiving mouse monoclonal antibodies for diagnosis or therapy. PSA levels may appear consistently elevated / depressed due to the interference by heterophilic antibodies & nonspecific protein binding. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be correlated with clinical findings and results of other investigations.

** End of Report **



Dr. Aarti Khanna Nagpal
DNB (Pathology)
Senior Consultant

