

Client
Gurugram
Pathkind Diagnostics Pvt. Ltd.
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

Processed By
Pathkind Diagnostics Pvt. Ltd.
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

Name : Mr. CL115	Billing Date : 07/07/2023 12:17:49
Age : 35 Yrs	Sample Collected on : 10/07/2023 10:01:31
Sex : Male	Sample Received on : 10/07/2023 11:02:13
P. ID No. : P1000100012597	Report Released on : 15/07/2023 16:56:39
Accession No : 10002304653	Barcode No. : 10002304653-01
Referring Doctor : Self	
Referred By :	Ref no. :

Report Status - Final

Test Name	Result	Biological Ref. Interval	Unit
BIOCHEMISTRY			
Testosterone Free & Total			
Testosterone Free <i>Sample: Serum</i> <i>Method: ELISA</i>	20.00	0 -21 Yrs- Not Defined 21 - 49 Yrs: 5.01 - 27.78 >50 Yrs: 4.11 - 21.85	pg/mL
Testosterone Total <i>Sample: Serum</i> <i>Method: ECLIA</i>	5.20	2.80 - 8.00	ng/mL

Testosterone Free

COMMENTS / INTERPRETATION :

- Free testosterone is a measure of biologically active testosterone, the value of which is unaffected by the variations in the transport proteins.
- The measurement of Free Testosterone is useful mainly in the evaluation of male hypogonadism and female hyperandrogenic states.
- Only 2-3 % of testosterone is unbound and free.

Testosterone Total

Clinical Significance :

Testosterone is the major androgenic hormone and is responsible for the development of the external genitalia and secondary sexual characteristics in males. It is an estrogen precursor in females, and in both genders, it has some anabolic effects and also influences behavior. High levels of testosterone during childhood leads to premature puberty in boys and masculinization in girls. Elevated levels in adult women results in varying degrees of virilization, including hirsutism, acne, oligo-amenorrhea and infertility. Mild-to-moderate testosterone elevations may be asymptomatic in males. Common causes of pronounced elevations of testosterone include congenital adrenal hyperplasia, adrenal, testicular, and ovarian tumors and abuse of testosterone or gonadotrophins by athletes. Low levels of testosterone is usually due to testicular failure in males, which can be primary, secondary or tertiary. It causes partial or complete hypogonadism and also causes some changes in the secondary sexual characteristics and the



Client
Gurugram
Pathkind Diagnostics Pvt. Ltd.
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

Processed By
Pathkind Diagnostics Pvt. Ltd.
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015


Name : Mr. CL115	Billing Date : 07/07/2023 12:17:49
Age : 35 Yrs	Sample Collected on : 10/07/2023 10:01:31
Sex : Male	Sample Received on : 10/07/2023 11:02:13
P. ID No. : P1000100012597	Report Released on : 15/07/2023 16:56:39
Accession No : 10002304653	Barcode No. : 10002304653-01
Referring Doctor : Self	
Referred By :	Ref no. :

Report Status - Final

Test Name	Result	Biological Ref. Interval	Unit
-----------	--------	--------------------------	------

reproductive function. In females, low levels of testosterone causes decline in libido and nonspecific mood changes.

** End of Report **



Dr. Saloni Garg
MD
Consultant Microbiology



Dr. Aarti Khanna Nagpal
DNB (Pathology)
Senior Consultant

