

**Client**  
**Gurugram**  
Pathkind Diagnostics Pvt. Ltd.  
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

**Processed By**  
**Pathkind Diagnostics Pvt. Ltd.**  
Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

<b>Name</b> : Mrs. SE173	<b>Billing Date</b> : 07/07/2023 12:35:13
<b>Age</b> : 26 Yrs	<b>Sample Collected on</b> : 10/07/2023 10:01:31
<b>Sex</b> : Female	<b>Sample Received on</b> : 10/07/2023 11:02:13
<b>P. ID No.</b> : P1000100013066	<b>Report Released on</b> : 19/07/2023 16:35:11
<b>Accession No</b> : 10002305122	<b>Barcode No.</b> : 10002305122-01
<b>Referring Doctor</b> : Self	<b>Ref no.</b> :
<b>Referred By</b> :	

**Report Status - Final**

Test Name	Result	Biological Ref. Interval	Unit
-----------	--------	--------------------------	------

**SEROLOGY**

<b># Mycoplasma Pneumoniae IgG Antibodies</b> <i>Sample: Serum</i> <i>Method: EIA</i>	5.24	Negative : < 9 Equivocal : 9 - 11 Positive : > 11	NTU
---	------	---	-----

**Mycoplasma Pneumoniae IgG Antibodies**

Interpretation

RESULT(INDEX)	REMARKS
<0.8	Negative
0.8-1.2	Equivocal
>1.2	Positive

Note

1. Positive IgM results are consistent with acute infection but false positives may occur
2. This assay should not be used as a screening procedure for the general population. Testing is recommended only if clinical evidence suggests the diagnosis of Mycoplasma pneumoniae associated disease.
3. Continued presence or absence of antibodies cannot be used to determine success or failure of therapy
4. Results should be interpreted in conjunction with clinical evaluation and other diagnostic procedures

Comments

Mycoplasma pneumoniae accounts for nearly 20% of all cases of pneumonia leading to Primary Atypical pneumonia. The disease has an insidious onset with fever and headache before the onset of respiratory symptoms. Symptomatic infections most commonly occur in children and young adults from 2-19 years. Negative IgM result does not rule out the presence of Mycoplasma pneumonia associated disease as the specimen may have been drawn before the appearance of detectable antibodies. If Mycoplasma infection is clinically indicated a second specimen should be submitted at least 14 days later.

\*\* End of Report \*\*

*Saloni Garg*

**Dr. Saloni Garg**

MD  
Consultant Microbiology

10002305122 Mrs. SE173

Page No: 1 of 1

