

**Client****Gurugram**

Pathkind Diagnostics Pvt. Ltd.

Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

**Processed By**

Pathkind Diagnostics Pvt. Ltd.

Plot No. 55-56, Udhog Vihar Ph-IV, Gurugram - 122015

<b>Name</b>	: Mrs. SE214	<b>Billing Date</b>	: 07/07/2023 12:36:19
<b>Age</b>	: 30 Yrs	<b>Sample Collected on</b>	: 10/07/2023 10:01:31
<b>Sex</b>	: Female	<b>Sample Received on</b>	: 10/07/2023 11:02:13
<b>P. ID No.</b>	: P1000100013109	<b>Report Released on</b>	: 17/07/2023 17:20:09
<b>Accession No</b>	: 10002305165	<b>Barcode No.</b>	: 10002305165-01
<b>Referring Doctor</b>	: Self		
<b>Referred By</b>	:	<b>Ref no.</b>	:

**Report Status - Final**

Test Name	Result	Biological Ref. Interval	Unit
-----------	--------	--------------------------	------

**SEROLOGY****Toxoplasma IgG Antibodies**

Sample: Serum  
Method: ECLIA

14.00 H

Negative : < 7.2  
Equivocal : 7.2 - 8.8  
Positive : > 8.8

IU/mL

**Toxoplasma IgG Antibodies****Interpretation:**

Negative : Presumed not to have had previous exposure or recent infection with Toxoplasma gondii.


Equivocal : May be due to low levels of IgG antibodies during the acute stage of infection. It is recommended to test those samples using a Toxoplasma IgM test, if clinically indicated.

Positive: Indicates past or recent infection with Toxoplasma gondii.

**Clinical Significance:**

- A negative result does not always exclude the possibility of Toxoplasma gondii infection.
- A single positive Toxoplasma IgG result should not be used to diagnose recent infection.
- Recent or acute infection can be elicited by testing for Anti Toxoplasma gondii IgM antibody.
- A suspected diagnosis of acute toxoplasmosis should be confirmed by detection of Toxoplasma gondii in cerebrospinal fluid or amniotic fluid by PCR.

\*\* End of Report \*\*

**Dr. Saloni Garg**

MD

Consultant Microbiology

